"I resisted starting insulin for years until my doctor recommended i-port Advance. Now, both she and I are very pleased with the results I've been able to achieve."

David, Age 52, Type 2

SIMPLE TO APPLY EASY TO USE. Use with standard syringe or pen Injection needle never pierces the skin Medication delivered Only a soft, flexible cannula under the skin

i-PORT ADVANCE™ IT TAKES THE **INJECTIONS FOR YOU** IT WON'T HURT **TO TRY**



A PAEDIATRIC DIABETES STUDY SHOWED THAT CHILDREN USING

IN JUST 6 MONTHS⁵

To order please contact Funding Management and ask for an i-Port Advance Account Set Up Form:

Call: **01923 205 167** (option 3)

Email: diabetesuk_fundingmgmt@medtronic.com

IMPORTANT SAFETY INFORMATION: i-Port Advance injection port is indicated for patients who administer, or to whom is administered, multiple daily subcutaneous injections of physician prescribed medications, including insulin. The device may remain in place for up to 72 hours to accommodate multiple drug injections without the discomfort of additional needle sticks. Site infection and/or irritation is a risk. Patients experiencing such symptoms should immediately consult their healthcare provider and/or discontinue use. For complete details, please consult i-Port Advance product user quide.

- 1 Hanas R, Adolfsson P, Elfvin-Akesson K et al. Indwelling catheters used from the onset of diabetes decrease injection pain and pre-injection anxiety. J Pediatr. 2002;140(3)315-320
- 2 Rabbone I, Bobbio A, Di Gianni V, Sacchetti C, Cerutti F. Intensive insulin therapy in preschool-aged diabetic children from multiple daily injections to continuous subcutaneous insulin infusion through indwelling catheters. J Endocrinol Invest. 2008;3 (3):193-195
- 3 Blevins T, Shwartz SL, Bode B et al. A study assessing an injection port for administration of insulin Diabetes Spectr. 2008:21(3):197-202
- 4 Riley D. Raup G. Impact of a subcutaneous injection device on improving patient care. Nurs Manage, 2010;41(6):49-50 5 HbA1C values in this study decreased from 9.4 to 8.5 in 6 months a 0.9% decrease. Burdick P, Cooper S,
- Horner B, Cobry E, McFann K, Chase HP. Use of a subcutaneous injection port to improve glycemic control in children with type 1 diabetes. Pediatr Diabetes, 2009:10:116-119
- 6 Rubin, Peyrot, Kruger, Travis. Barriers to Insulin Injection Therapy: Patient and Health Care Provider Perspectives The Diabetes Educator 2009; 35;1014

United Kingdom Medtronic Limited **Building 9, Croxley Park** Hatters Lane Watford, Herts WD188WW Tel: +44 (0)1923 205 167 Fax: +44 (0)1923 241 004 www.medtronic-diabetes.co.uk

Medtronic Ireland Limited Block 3090-3094

Lake Drive

Citywest Business Campus Dublin **DN24 XN47**

Tel: +353 (0)15111499 Fax: +353 (0)18077220 www.medtronic-diabetes.ie

Medtronic



LIC201705345 FN





Medtronic

3 DAYS **PUNCTURE FREE**

After the application of your i-Port Advance you can easily administer multiple injections!

Over the course of 3 days, you will take up to 15 injections. With **i-Port Advance** it's only a single application.

LESS PAIN¹

LESS ANXIETY¹



IMPROVES QUALITY OF LIFE²

DO YOU ALTER ACTIVITIES TO **AVOID INJECTIONS?**

43% ALTER **EATING HABITS**

19% **PHYSICAL ACTIVITY**⁶ Continue your current diabetes treatment with one i-Port Advance³





BASAL e.g., insulin glargine, insulin detemir


i-Port Advance can be used for Basal and Bolus Doses

60 MIN



15% **ALTER SOCIAL EVENTS**⁶

WITH INJECT ONCE EVERY 3 DAYS VERSUS **EVERY DAY WITH STANDARD INJECTIONS**



100% REPORTED THAT i-PORT ADVANCE HELPED THEM TO **DECREASE ANXIETY**⁴

99% OF PATIENTS REPORTED THAT i-PORT ADVANCE WAS BENEFICIAL⁴